

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|---|
| (51) International Patent Classification ⁶ : C12N 9/10, 9/24, 9/42 | A1 | (11) International Publication Number: WO 98/38288 (43) International Publication Date: 3 September 1998 (03.09.98) |
| (21) International Application Number: PCT/DK98/00076 (22) International Filing Date: 26 February 1998 (26.02.98) (30) Priority Data: 0212/97 26 February 1997 (26.02.97) DK 60/051,016 27 May 1997 (27.05.97) US (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventor; and (75) Inventor/Applicant (for US only): NIELSEN, Ruby, Illum [DK/DK]; Gedebakken 9, DK-3520 Farum (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK). | | (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: MICROBIAL XYLOGLUCAN ENDOTRANSGLYCOSYLASE (XET) (57) Abstract It has been found by a screening assay that XET activity is produced by an overwhelming array of phylogenetically dispersed microorganisms. Accordingly, the present invention relates to a xyloglucan endotransglycosylase preparation which is producible by cultivation of a microorganism expressing an XET. | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

5 **MICROBIAL XYLOGLUCAN ENDOTRANSGLYCOSYLASE (XET)****FIELD OF THE INVENTION**

The present invention relates to microbial xyloglucan endotransglycosylases (XETs) and the production and uses thereof.

10

BACKGROUND OF THE INVENTION

Xyloglucan endotransglycosylase (XET) is an enzyme known from plants. To our best knowledge XET has never been described from microorganisms.

15

Stephen C. Fry et al. suggest in Biochem. J (1992) 282, p. 821-828 that XET is responsible for cutting and rejoining intermicrofibrillar xyloglucan chains and that XET thus causes the wall-loosening required for plant cell expansion.

20

XET has been suggested for use in regulating the morphology of a plant, see EP 562 836 p. 2 l. 27-28.

25

XET is believed to be present in all plants, in particular in all land plants. XET has been extracted from dicotyledons, monocotyledons, in particular graminaceous monocotyledons and liliaceous monocotyledons, and also from a moss and a liverwort, for reference see Biochem. J (1992) 282, p. 823.

30

In the copending patent application PCT/DK 96/00538 (WO 97/23683) we have shown that a cellulosic material may get improved strength properties and/or improved shape-retention properties and/or improved anti-wrinkling properties after treatment with an XET enzyme.

35

The XET enzyme has some very interesting applications, so it is an object of the present invention to provide microbially derived xyloglucan endotransglycosylases useful for, e.g., the applications described above. The microbially derived xyloglucan endotransglycosylases would have the great advantage that they could easily be produced in great quantities.

BRIEF DISCLOSURE OF THE INVENTION

40

5 It has been found by a screening assay that XET activity is produced by an overwhelming array of phylogenetically dispersed microorganisms.

 Accordingly, the present invention relates to a xyloglucan
10 endotransglycosylase preparation which is producible by cultivation of a microorganism expressing an XET; in particular the present invention relates to:

A method for the production of a xyloglucan endotransglycosylase
15 enzyme (XET) comprising

- (a) culturing in a suitable nutrient medium a microorganism expressing a microbial XET under conditions conducive to the production of the XET enzyme, and
- (b) subsequently recovering of the XET enzyme from the nutrient
20 medium.

 Further, the present invention relates to use of the XET preparation of the present invention for treating cellulosic material, and to microbial XET preparations as such.

25

BRIEF DESCRIPTION OF DRAWINGS

 The present invention is further illustrated by reference to the accompanying drawing, in which

30 Fig. 1 shows the pH profiles of *Dichotomocladium hesseltinei*, *Tiarosporella phaseolina* and *Pseudoplectania nigrella* XET activity (Δ : *Tiarosporella phaseolina*; \square : *Pseudoplectania nigrella*; and \blacklozenge : *Dichotomocladium hesseltinei*) obtained according to Example 5.

35 Fig. 2 shows the pH profiles of *Dichotomocladium hesseltinei*, *Tiarosporella phaseolina* and *Pseudoplectania nigrella* xyloglucanase activity (Δ : *Tiarosporella phaseolina*; \square : *Pseudoplectania nigrella*; and \blacklozenge : *Dichotomocladium hesseltinei*) obtained according to Example 5.

5

DETAILED DISCLOSURE OF THE INVENTION

The present invention relates to a xyloglucan endotransglycosylase preparation which is producible by cultivation of a microorganism.

It has been shown in the present application that XET is produced by fungi and bacteria, and it is contemplated that XET is also produced by yeasts.

An "XET-paper" was used for identifying XET producing microorganisms. The "XET-paper" is described in the copending patent application PCT/GB96/02351 (WO 97/11193); it is a xyloglucan-coated paper which is dipped through a labelled oligosaccharide.

The labelled oligosaccharide, the xyloglucan-impregnated paper, and the dot-blot test for XET activity were made in the following way:

Preparation of labelled oligosaccharide

The reducing oligosaccharide 4-O-[4-O-[4-O-[6-O- α -D-Xylopyranosyl- β -D-glucopyranosyl]-6-O-(2-O- β -D-galactopyranosyl)- α -D-xylopyranosyl- β -D-glucopyranosyl]-6-O-(2-O- β -D-galactopyranosyl)- α -D-xylopyranosyl- β -D-glucopyranosyl]-D-glucose ("XLLG") (1 gram) is dissolved in 25 ml of a saturated aqueous solution of ammonium hydrogencarbonate containing 1 gram of sodium cyanoborohydride (NaCNBH₃) and incubated in the dark at 25°C for 7 days to permit reductive amination. The ammonium hydrogencarbonate is then removed by drying, and the (ninhydrin-reactive) aminated derivative of XLLG is purified e.g. by gel-permeation chromatography or cation-exchange chromatography. The product is believed to be an oligosaccharidyl-1-amino-1-deoxyalditol, i.e. a derivative of XLLG in which the reducing terminal D-glucose moiety has been replaced by 1-amino-1-deoxy-D-glucitol.

5 The oligosaccharidyl-1-amino-1-deoxyalditol (50mg) is dissolved in 3 ml of 3% borax (di-sodium tetraborate; pH = 9.0-9.5) and a freshly-prepared solution of 10 mg lissamine rhodamine sulphonyl chloride [purchased from Molecular Probes Inc., USA] in 0.75 ml of dry dimethylformamide (DMF) is added gradually, with
10 stirring, and the mixture is incubated in the dark overnight. A further 0.75 ml of DMF containing 10 mg lissamine rhodamine sulphonyl chloride is added and the mixture incubated for a further 8 h. The bright pink oligosaccharidyl-1-amino-1-deoxyalditol-lissamine-rhodamine conjugate (XLLGol-SR) is
15 purified by gel-permeation chromatography followed by reversed-phase chromatography on a C_{18} -silica gel column. After washing of the latter column with water, a methanol gradient is applied and the XLLGol-SR elutes in about 50% methanol.

20 Preparation of xyloglucan-impregnated paper

 Whatman No.1 filter paper is moistened with a 1% aqueous solution of xyloglucan and dried. The XLLGol-SR preparation is diluted into enough 75% aqueous acetone to give an absorbance at
25 580 nm (A_{580}) of 0.2; the xyloglucan-coated sheet of Whatman No. 1 paper is then dipped through this solution and re-dried; the product is referred to as "XET-paper". Suitably sized pieces (e.g. 72 x 108 mm) of the XET-paper may then be glued with a non-aqueous adhesive onto a non-absorbent medium such as a sheet of
30 transparent acetate.

Dot-blot test for XET activity

35 (i) A spot of the solution to be tested for XET-activity is pipetted on to a marked position in a piece of XET-paper. If the spots are 4 μ l, the spacing between the samples can conveniently be 9 mm (centre-to-centre, i.e. as in a standard 96-well test plate format).

5 (ii) The XET paper is then quickly (before the spots have dried) clamped between two sheets of plastic (e.g. acetate sheets, as used on overhead projectors) and incubated e.g. at 20°C for 1 hour.

10 (iii) The incubated XET-paper and its plastic backing is then placed (paper-side down) in a dish containing about 150 ml of a solvent [e.g. freshly prepared ethanol/formic acid/water (1:1:1 by volume)] that will remove from the paper the unreacted XLLGol-SR but not any XLLGol-SR that has become incorporated into
15 the xyloglucan owing to XET-catalysed transglycosylation. The paper now readily detaches from the plastic backing.

(iv) The paper is then rinsed in running water for 5
20 minutes, then in approximately 100 ml of acetone for 5 minutes, and then dried thoroughly. If desired, drying can be expedited by a 5-minute treatment in an oven at 80°C.

(v) The paper is then examined under a short-wavelength
25 ultraviolet lamp (e.g. emitting at 254 nm; suitable eye- and skin-protection should be worn). Active XET is indicated by a pink (orange-fluorescing) spot, which can be quantified, e.g. by use of a scanning spectrofluorimeter.

30 XET enzymes

We have discovered that microbial enzymes with XET activity may be either transglycosylases (meaning that they lack hydrolase activity or only have a very low hydrolase activity) or they may catalyze both transglycosylation and hydrolysis of xyloglucan.

35

XET enzymes with both transglycosylating and hydrolytic activities are also described for XET enzymes obtained from plants (see Annu. Rev. Plant Physiol. Plant Mol. Biol. 1995.46: p. 509).

40

5 Taxonomic classification

The taxonomic classification used herein builds primarily on the system used in the :NIH Data Base (Entrez, version spring 1996) available on World Wide Web: (<http://www3.ncbi.nlm.nih.gov/htbin/ef/entrezTAX>).

10 Regarding classification of organisms which are not included in the Entrez data base the following generally available and world wide accepted reference books have been used:

For Ascomycetes: Eriksson, O.E. & Hawksworth, D.L.: Systema Ascomycetum vol 12 (1993).

For Basidiomycetes, Jülich, W.: Higher Taxa of Basidiomycetes, Bibliotheca Mycologia 85, 485pp (1981).

For Zygomycetes: O'Donnell, K.: Zygomycetes in culture, University of Georgia, US, 257pp (1979).

20 General mycological reference books:

Hawksworth, D.L., Kirk, P.M., Sutton, B.C. and Pegler, D.N.: Dictionary of the fungi, International Mycological Institute, 616 pp (1995);

25 Von Arx, J. A.: The genera of fungi sporulating in culture, 424 pp (1981).

Preferred fungi

In a preferred embodiment of the present invention the XET preparation is produced by cultivation of a fungus, in particular
30 a fungus which belongs to Basidiomycota, Zygomycota, Ascomycota or a Mitosporic fungus.

A preferred Basidiomycota strain is a Hymenomycetes strain belonging to the orders Coriolales, Schizophyllales, Stereales or Xenasmatales; in particular a strain belonging to one of the
35 families Coriolaceae, Corticiaceae, Schizophyllaceae, Stereaceae or Tubulicrinaceae. Preferred genera is one of the following: Trametes, Corticium, Schizophyllum, or Tubulicrinis. Preferred species is one of the following Trametes hirsuta, Corticium roseum, Schizophyllum sp, Stereum hirsutum or Tubulicrinis

5 subulatus.

Preferred Ascomycota are strains belonging to the classes Loculoascomycetes, Discomycetes, Pyrenomycetes, and Plectomycetes, preferably those belonging to the orders Dothideales, Rhytismatales, Pezizales, Leotiales, Xylariales, Hypocreales, Halosphaeriales, Phyllachorales, Diaporthales and Eurotiales.

Preferred strains are strains belonging to the families Botryosphaeriaceae, Dothioraceae, Mycosphaerellaceae, Tubeufiaceae, Pleosporaceae, Leptosphaeriaceae, Rhytismataceae, Sarcosomataceae, Pyronemataceae, Ascobolaceae, Sclerotiniaceae, Amphisphaeriaceae, Xylariaceae, Hypocreaceae, Halosphaeriaceae, Phyllachoraceae, Valsaceae, Melanconidaceae and Trichocomataceae; especially strains belonging to the genera Diplodia, Plowrightia, Phyllosticta, Septoria, Tubeufia, Alternaria, Coniothyrium, Phoma, Embellisia, Tiarosporella, Galiella, Pseudoplectania, Pyronema, Oedocephalum, Botrytis, Aposphaeria, Pestalotia, Pestalotiopsis, Poronia, Nodulisporium, Xylaria, Fusarium, Verticillium, Volutella, Chaetapiospora, Lulworthia, Colletotrichum, Cytospora, Discula, Phomopsis, Coryneum, Seimatosporium, Aspergillus, Eurotium, Eupenicillium, Penicillium, Petromyces and Talaromyces.

Preferred are the species Diplodia gossypina, Plowrightia ribesia, Phyllosticta sp, Septoria sp, Tubeufia amazonensis, Alternaria sp, Embellisia hyacinthi, Phoma neoloba, Phoma tropica, Coniothyrium sp, Coniothyrium olivaceum, Coniothyrium dunckii, Tiarosporella phaseolina, Tiarosporella sp, Galiella celebica, Pseudoplectania nigrella, Pyronema domesticum, Oedocephalum sp, Botrytis cinerea, Aposphaeria sp, Pestalotia sp, Pestalotiopsis sp, Poronia punctata, Xylaria sp, Nodulisporium sp, Fusarium solani, Verticillium sp, Volutella buxi, Chaetapiospora rhododendri, Lulworthia uniseptata, Colletotrichum aculatum, Colletotrichum crassipes, Cytospora spp,

5 *Discula* sp, *Phomopsis ilicis*, *Phomopsis cirsii*, *Coryneum*
castaneicola, *Seimatosporium lichenicola*, *Aspergillus tamarii*,
Eurotium chevalieri, *Eupenicillium javanicum*, *Penicillium*
capsulatum, *Penicillium olsonii*, *Penicillium pinophilum*,
Penicillium roqueforti, *Penicillium italicum*, *Penicillium*
10 *canescens*, *Penicillium verruculosum*, *Petromyces alliaceus* and
Talaromyces flavus.

Examples of useful Zygomycota are strains belonging to the order Mucorales, preferably strains belonging to the families Chaetocladiaceae and Mucoraceae.

15 Preferred strains belong to the genera *Dichotomocladium*, *Actinomucor*, *Gongronella*, *Sporodiniella* and *Mucor*, in particular *Dichotomocladium hesseltinei*, *Actinomucor elegans*, *Gongronella butleri*, *Sporodiniella umbellata* and *Mucor miehei* var *minor*.

20 Example of a strain of uncertain taxonomy is *Vialaea insculpta*.

Examples of strains belonging to the Mitosporic
25 Fungi are *Acrodontium crateriforme*, *Aureobasidium pullulans*, *Circinotrichum* sp, *Cryptocline* sp, *Ellisiopsis* sp, *Epicoccum nigrum*, *Gliocladium* sp, *Helicorhoidium irregulare*, *Hendersonia* spp, *Mariannaea* sp, *Microsphaeropsis* sp, *Ramularia* sp, *Sarcopodium* sp, *Spadicoides* sp, *Speiropsis pedatospora*,
30 *Sporotrichum exile*, *Stilbella* sp, *Trichothecium* sp, *Trimmatostroma abietes*, *Tubakia dryina*, *Wiesneriomyces* sp and *Zygosporium masonii*.

Preferred bacteria

35 In another aspect, the invention relates to a novel XET preparation which is producible by cultivation of a bacterium.

Preferred bacteria are gram-negative or gram-positive.

5 Examples of XET-producing gram-positive bacteria are strains belonging to the genus *Bacillus*.

Preferred strains

10 The following strains have been found to be XET-positive:

1. *Dichotomocladium hesseltinei*. Acc No of strain: CBS 164.61.
Classification: Zygomycota, Mucorales, Chaetocladiaceae.
2. *Actinomucor elegans*. Ex of Acc No of strain: CBS 154.86.
15 Classification: Zygomycota, Mucorales, Chaetocladiaceae.
3. *Mucor miehei* var *minor*. Acc No of strain: ATCC 36018.
Classification: Zygomycota, Mucorales, Mucoraceae.
4. *Gongronella butleri*. A strain of *Gongronella butleri* has been deposited according to the Budapest Treaty on the International
20 Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 448.97.
Classification: Zygomycota, Mucorales, Chaetocladiaceae.
5. *Sporodiniella umbellata*. Acc No of strain: CBS 195.77.
25 Classification: Zygomycota, Mucorales, Mucoraceae.
6. *Phyllosticta* sp. Isolated from a leaf of *Pithecolobium* sp from China. Classification: Ascomycota, Loculoascomycetes, Dothidiales, Mycosphaellaceae.
7. *Septoria* sp. A strain of *Septoria* sp has been deposited
30 according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 2 January 1996, at Centraalbureau voor Schimmelcultures (CBS), under Accession No. CBS 831.95.

- 5 Classification: *Ascomycota*, *Loculoascomycetes*, *Dothidiales*,
Mycosphaellaceae.
8. *Diplodia gossypina*. A strain of *Diplodia gossypina* has been
deposited according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of
10 Patent Procedures, on 12 March 1996, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 274.96.
Classification: *Ascomycota*, *Loculoascomycetes*, *Dothidiales*,
Botryosphaeriaceae.
9. *Plowrightia ribesia*. Isolated from *Ribes* sp., Denmark.
15 Classification: *Ascomycota*, *Loculoascomycetes*, *Dothidiales*,
Dothioraceae.
10. *Tubeufia amazonensis*. Acc No of Strain: ATCC 42524.
Classification: *Ascomycota*, *Loculoascomycetes*, *Dothidiales*,
Tubeufiaceae.
- 20 11. *Alternaria* sp. Classification: *Ascomycota*,
Loculoascomycetes, *Dothidiales*, *Pleosporaceae*.
12. *Embellisia hyacinthi*. Acc No of species IMI 211561.
Classification: *Ascomycota*, *Loculoascomycetes*, *Dothidiales*,
Pleosporaceae.
- 25 13. *Phoma neoloba*. Classification: *Ascomycota*,
Loculoascomycetes, *Dothideales*, *Pleosporaceae*.
14. *Phoma tropica*. Ex on Acc No of species CBS 537.66.
Classification: *Ascomycota*, *Loculoascomycetes*, *Dothideales*,
Pleosporaceae.
- 30 15. *Coniothyrium* sp. Classification: *Ascomycota*,
Loculoascomycetes, *Dothideales*, *Leptosphaeriaceae*.
16. *Coniothyrium olivaceum*. Ex on Acc No of species CBS 304.68.
Classification: *Ascomycota*, *Loculoascomycetes*, *Dothideales*,

5 *Leptosphaeriaceae*.

17. *Coniothyrium duncckii*. Classification: Ascomycota, Loculoascomycetes, Dothideales, Leptosphaeriaceae.

18. *Tiarosporella phaseolina*. A strain of *Tiarosporella phaseolina* (*Macrophomina* sp) has been deposited according to the
10 Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on the 28th January 1997, at Centraalbureau voor Schimmelcultures (CBS) under Accession No. CBS 446.97. Classification: Ascomycota, Discomycetes, Rhytismatales, Rhytismataceae.

15 19. *Tiarosporella* sp. Classification: Ascomycota, Discomycetes, Rhytismatales, Rhytismataceae.

20. *Galiella celebica*. Isolated from a sample collected in Japan. Classification: Ascomycota, Discomycetes, Pezizales, Sarcosomataceae.

20 21. *Pseudoplectania nigrella*. A strain of *Pseudoplectania nigrella* has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 28 January 1997, at Centraalbureau voor Schimmelcultures (CBS), under Accession No.
25 CBS 444.97. Classification: Ascomycota, Discomycetes, Pezizales, Sarcosomataceae.

22. *Pyronema domesticum*. Isolated from a sample from Norway. Classification: Ascomycota, Discomycetes, Pezizales, Pyronemataceae.

30 23. *Oedocephalum* sp. Classification: Ascomycota, Discomycetes, Pezizales, Ascobolaceae.

24. *Botrytis cinerea*. A strain of *Botrytis cinerea* has been deposited according to the Budapest Treaty on the International

- 5 Recognition of the Deposits of Microorganisms for the Purpose of
Patent Procedures, on 28 January 1997, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 447.97.
Classification: Ascomycota, Discomycetes, Leotiales,
Sclerotiniaceae.
- 10 25. *Aposphaeria* sp. Classification: Ascomycota, Discomycetes,
Leotiales, Sclerotiniaceae.
26. *Pestalotia* sp. A strain of *Pestalotia* sp. has been deposited
according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of
15 Patent Procedures, on 28 January 1997, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 445.97.
Classification: Ascomycota, Pyrenomycetes, Xylariales,
Amphisphaeriaceae.
27. *Pestalotiopsis* sp. Classification: Ascomycota,
20 Pyrenomycetes, Xylariales, Amphisphaeriaceae.
28. *Poronia punctata*. Isolated from a sample from Sweden.
Classification: Ascomycota, Pyrenomycetes, Xylariales,
Xylariaceae.
29. *Xylaria* sp. Isolated from a leaf of the palm, *Sabal*
25 *jamaicensis*, growing in Mona, Jamaica. Classification:
Ascomycota, Pyrenomycetes, Xylariales, Xylariaceae Xylariaceae.
30. *Nodulisporium* sp. Classification: Ascomycota, Pyrenomycetes,
Xylariales, Xylariaceae.
31. *Fusarium solani*. Isolated from a sample of grain of maize
30 from India. Classification: Ascomycota, Pyrenomycetes,
Hypocreales, Hypocreaceae.
32. *Verticillium* sp. A strain of *Verticillium* sp has been
deposited according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of

- 5 Patent Procedures, on 2 January 1996, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 830.95.
Classification: Ascomycota, Pyrenomycetes, Hypocreales,
Hypocreaceae.
33. *Volutella buxi*. Acc No of Strain: IMI 049467.
10 Classification: Ascomycota, Pyrenomycetes, Hypocreales,
Hypocreaceae.
34. *Chaetapiozpora rhododendri*. Classification: Ascomycota,
Pyrenomycetes, Xylariales, Xylariaceae, Hyponectriaceae.
35. *Lulworthia uniseptata*. A strain of *Lulworthia uniseptata* has
15 been deposited according to the Budapest Treaty on the
International Recognition of the Deposits of Microorganisms for
the Purpose of Patent Procedures, on 28 January 1997, at
Centraalbureau voor Schimmelcultures (CBS), under Accession No.
CBS 442.97. Classification: Ascomycota, Pyrenomycetes,
20 Halosphaeriales, Halosphaeriaceae.
36. *Colletotrichum aculatum*. Classification: Ascomycota,
Pyrenomycetes, Phyllachorales, Phyllachoraceae.
37. *Colletotrichum crassipes*. Classification: Ascomycota,
Pyrenomycetes, Phyllachorales, Phyllachoraceae.
- 25 38. *Cytospora* sp. A strain of *Cytospora* sp has been deposited
according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of
Patent Procedures, on 23 January 1997, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 424.97.
30 Classification: Ascomycota, Pyrenomycetes, Diaporthales,
Valsaceae.
39. *Cytospora* sp. A strain of *Cytospora* sp has been deposited
according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of
35 Patent Procedures, on 23 January 1997, at Centraalbureau voor

- 5 Schimmelcultures (CBS), under Accession No. CBS 425.97.
Classification: Ascomycota, Pyrenomycetes, Diaporthales,
Valsaceae.
40. *Discula* sp. Classification: Ascomycota, Pyrenomycetes,
Diaporthales, Valsaceae.
- 10 41. *Phomopsis ilicis*. Classification: Ascomycota, Pyrenomycetes,
Diaporthales, Valsaceae.
42. *Phomopsis cirsii*. Classification: Ascomycota, Pyrenomycetes,
Diaporthales, Valsaceae.
43. *Coryneum castaneicola*. Classification: Ascomycota,
15 Pyrenomycetes, Diaporthales, Melanconidaceae.
44. *Seimatosporium lichenicola*. Classification: Ascomycota,
Pyrenomycetes, Diaporthales, Melanconidaceae.
45. *Aspergillus tamarii*. Ex of Acc No of strain: CBS 821.72.
Classification: Ascomycota, Plectomycetes, Eurotiales,
20 Trichocomataceae.
46. *Eurotium chevalieri*. Ex of Acc No of strain: CBS 472.91.
Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
47. *Penicillium capsulatum*. Ex of Acc No of strain: CBS 273.86.
25 Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
48. *Penicillium olsonii*. Ex of Acc No of strain: CBS 523.89.
Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
- 30 49. *Penicillium pinophilum*. Ex of Acc No of strain: CBS 440.89.
Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.

- 5 50. *Penicillium roqueforti*. Ex of Acc No of strain: CBS 167.91.
Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
51. *Penicillium italicum*. Ex of Acc No of strain: IMI 078 681.
Classification: Ascomycota, Plectomycetes, Eurotiales,
10 Trichocomataceae.
52. *Penicillium canescens*. Ex of Acc No of strain: CBS 579.70.
Isolated from a salt mine in Egypt. Classification: Ascomycota,
Plectomycetes, Eurotiales, Trichocomataceae.
53. *Eupenicillium javanicum*. Ex of Acc No of the strain: CBS
15 448.74. Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
54. *Penicillium verruculosum*. Ex of Acc No of strain: CBS
563.92. Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
- 20 55. *Talaromyces flavus*. Acc No of the strain: ATCC 52201.
Classification: Ascomycota, Plectomycetes, Eurotiales,
Trichocomataceae.
56. *Petromyces alliaceus*. Acc No of strain: CBS 511.69.
Classification: Ascomycota, Plectomycetes, Eurotiales,
25 Trichocomataceae.
57. *Trametes hirsuta*. Isolated from a sample collected in
Denmark. Classification: Basidiomycota, Hymenomycetes,
Coriolales, Coriolaceae.
58. *Schizophyllum* sp. A strain of *Schizophyllum* sp has been
30 deposited according to the Budapest Treaty on the International
Recognition of the Deposits of Microorganisms for the Purpose of
Patent Procedures, on 28 January 1997, at Centraalbureau voor
Schimmelcultures (CBS), under Accession No. CBS 443.97.

- 5 Classification: *Basidiomycota*, *Hymenomycetes*, *Schizophyllales*,
Schizophyllaceae.
59. *Corticium roseum*. Isolated from a sample collected in
Denmark. Classification: *Basidiomycota*, *Hymenomycetes*,
Aleurodiscales Cortiaceae.
- 10 60. *Tubulicrinis subulatus*. Isolated from a sample collected in
Denmark. Classification: *Basidiomycota*, *Hymenomycetes*,
Xenasmatales, Tubulicrinaceae.
61. *Stereum hirsutum*. Isolated from a sample collected in
Denmark. Classification: *Basidiomycota*, *Hymenomycetes*,
15 *Stereales, Stereaceae*.
62. *Acrodontium crateriforme*. Classification: *Mitosporic fungus*.
63. *Aureobasidium pullulans*. Classification: *Mitosporic fungus*.
64. *Circinotrichum* sp. Classification: *Mitosporic fungus*.
65. *Cryptocline* sp. Classification: *Mitosporic fungus*.
- 20 66. *Ellisiopsis* sp. Classification: *Mitosporic fungus*.
67. *Epicoccum nigrum*. Classification: *Mitosporic fungus*.
68. *Gliocladium* sp. Classification: *Mitosporic fungus*.
69. *Helicorhoidion irregulare*. Classification: *Mitosporic
fungus*.
- 25 70. *Hendersonia* sp. Classification: *Mitosporic fungus*.
71. *Mariannaea* sp. Classification: *Mitosporic fungus*.
72. *Microsphaeropsis* sp. Classification: *Mitosporic fungus*.
73. *Ramularia* sp. Classification: *Mitosporic fungus*.
74. *Sarcopodium* sp. Classification: *Mitosporic fungus*.
- 30 75. *Spadicoides sto*. Acc No of strain IMI203428. Classification:
Mitosporic fungus.
76. *Speiropsis pedatospora*. Classification: *Mitosporic fungus*.
77. *Sporotrichum exile*. Acc No of strain CBS 350.47.
Classification: *Mitosporic fungus*.
- 35 78. *Stilbella* sp. Classification: *Mitosporic fungus*.

- 5 79. *Trichothecium* sp. Classification: Mitosporic fungus.
80. *Trimmatostroma abietes*. Classification: Mitosporic fungus.
81. *Tubakia dryina*. Classification: Mitosporic fungus.
82. *Wiesneriomyces* sp. Classification: Mitosporic fungus.
83. *Zygosporium masonii*. Classification: Mitosporic fungus.
- 10 84. *Vialaea insculpta*. Classification: Uncertain.
85. *Bacillus alcalophilus*. A strain of *Bacillus alcalophilus* has been deposited according to the Budapest Treaty on the International Recognition of the Deposits of Microorganisms for the Purpose of Patent Procedures, on 12 February 1997, at
15 Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, under Accession No. DSM 11404.

Production of XET

20 The XET enzyme of the invention may be produced by aerobic cultivation of the above mentioned microbial strains on a nutrient medium containing suitable carbon and nitrogen sources, such media being known in the art.

Alternatively, the XET enzyme of the invention may be produced by aerobic cultivation of a transformed host organism
25 containing the appropriate genetic information from, e.g., one of the above mentioned strains. Accordingly, the present invention also relates to a method for the production of a xyloglucan endotransglycosylase enzyme (XET) comprising

- (a) culturing in a suitable nutrient medium a transformed host
30 microorganism expressing a microbial XET under conditions conducive to the production of the XET enzyme, and
(b) subsequently recovering of the XET enzyme from the nutrient medium.

Such transformants can be prepared and cultivated by
35 methods known in the art:

Cloning a DNA Sequence Encoding XET

The DNA sequence encoding a XET enzyme of the

5 invention may be isolated from any microorganism producing the XET in question, using various methods well known in the art.

First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the XET to be studied. Then, if the amino
10 acid sequence of the XET is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify XET-encoding clones from a genomic library or cDNA prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known XET gene could
15 be used as a probe to identify XET-encoding clones, using hybridization and washing conditions of lower stringency.

Yet another method for identifying XET-encoding clones would involve inserting fragments of genomic DNA or cDNA into an expression vector, such as a plasmid, transforming XET-negative
20 host organism with the resulting DNA library, then plating the transformed cells onto agar plates and by use of the assay described above allowing clones expressing the XET to be identified.

Alternatively, the DNA sequence encoding the enzyme
25 may be prepared synthetically by established standard methods, e.g. the phosphoamidite method described by S.L. Beaucage and M.H. Caruthers in Tetrahedron Letters 22, 1981, pp. 1859-1869 or the method described by Matthes et al. in The EMBO J. 3, 1984, pp. 801-805. In the phosphoamidite method, oligonucleotides are
30 synthesized, e.g., in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of
35 synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in US 4,683,202 or R.K. Saiki
40 et al. in Science 239, 1988, pp. 487-491.

5

Expression of XET

According to the invention, a XET-encoding DNA sequence produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding a XET enzyme of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding a XET of the invention, especially in a bacterial host, are the promoter of the *lac* operon of *E.coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis* α -amylase gene (*amyL*), the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus Amyloliquefaciens* α -amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host,

5 examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral α -amylase, *A. niger* acid stable α -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or
10 *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the XET enzyme of the invention. Termination and poly-
15 adenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of
20 plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g., a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one which confers antibiotic resistance such as
25 ampicillin, kanamycin, chloramphenicol or tetracyclin resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to hygromycin resistance, or the selection may be accomplished by co-transformation, e.g., as described in WO 91/17243.

30 While intracellular expression may be advantageous in some respects, e.g., when using certain bacteria as host cells, it is generally preferred that the expression is extracellular.

Procedures suitable for constructing vectors of the invention encoding a XET enzyme and containing the promoter, terminator and other elements, respectively, are well known to
35 persons skilled in the art (cf., for instance, Sambrook et al. in Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989).

The cell of the invention, either comprising a DNA

5 construct or an expression vector of the invention as defined
above, is advantageously used as a host cell in the recombinant
production of a XET of the invention. The cell may be transformed
with the DNA construct of the invention encoding the XET conveni-
10 the host chromosome. This integration is generally considered to
be an advantage as the DNA sequence is more likely to be stably
maintained in the cell. Integration of the DNA constructs into
the host chromosome may be performed according to conventional
methods, e.g., by homologous or heterologous recombination.
15 Alternatively, the cell may be transformed with an expression
vector as described above in connection with the different types
of host cells.

The cell of the invention may be a cell of a higher
organism such as a mammal or an insect, but is preferably a
20 microbial cell, e.g., a bacterial or a fungal (including yeast)
cell.

Examples of suitable bacteria are grampositive
bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*,
Bacillus lentus, *Bacillus brevis*, *Bacillus stearothermophilus*,
25 *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus*
coagulans, *Bacillus circulans*, *Bacillus lautus*, *Bacillus*
megaterium, *Bacillus thuringiensis*, or *Streptomyces lividans* or
Streptomyces murinus, or gramnegative bacteria such as *E.coli*.
The transformation of the bacteria may, for instance, be effected
30 by protoplast transformation or by using competent cells in a
manner known per se.

The yeast organism may favourably be selected from a
species of *Saccharomyces* or *Schizosaccharomyces*, e.g., *Sacchar-*
omyces cerevisiae. The filamentous fungus may advantageously
35 belong to a species of *Aspergillus*, e.g., *Aspergillus oryzae* or
Aspergillus niger. Fungal cells may be transformed by a process
involving protoplast formation and transformation of the
protoplasts followed by regeneration of the cell wall in a manner
known per se. A suitable procedure for transformation of

5 *Aspergillus* host cells is described in EP 238 023.

In a yet further aspect, the present invention relates to a method of producing a XET enzyme of the invention, which method comprises cultivating a host cell as described above under conditions conducive to the production of the XET and recovering
10 the XET from the cells and/or culture medium.

The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the XET of the invention. Suitable media are available from commercial suppliers or may be
15 prepared according to published recipes (e.g., as described in catalogues of the American Type Culture Collection).

The XET secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells from the medium by centrifugation
20 or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

25 Industrial Applications

According to the present invention a cellulosic material may get improved strength properties and/or improved shape-retention properties and/or improved anti-wrinkling properties after treatment with a XET enzyme. The XET enzyme
30 has the ability to rearrange and link the xyloglucan molecules which are hydrogen bonded to the cellulosic fibres whereby the above mentioned features may be achieved.

In order to enhance the effect of the XET enzyme it may in some cases be an advantage to add xyloglucan to the cellulosic
35 material whereby the enzyme may be able to link more cellulosic material together.

The treatment of the cellulosic material with the XET enzyme may be carried out in water, or it may be carried out in water in the presence of certain components such as a buffer
40 and/or a wetting agent and/or a stabilizer and/or a polymer

5 and/or an organic component reducing the water activity such as DMSO.

The buffer may suitably be a phosphate, borate, citrate, acetate, adipate, triethanolamine, monoethanolamine, diethanolamine, carbonate (especially alkali metal or alkaline
10 earth metal, in particular sodium or potassium carbonate, or ammonium and HCl salts), diamine, especially diaminoethane, imidazole, Tris or amino acid buffer.

The wetting agent serves to improve the wettability of the cellulosic material. The wetting agent is preferably of a
15 non-ionic surfactant type.

The stabilizer may be an agent stabilizing the XET enzyme.

According to the invention the concentration of XET in the aqueous medium may be 0.01-1000 µg of enzyme protein per g
20 cellulosic material, preferably 0.05-100 µg of enzyme protein per g cellulosic material.

It will generally be appropriate to incubate the reaction medium (containing the cellulosic material and the XET enzyme) for a period of at least a few minutes. An incubation
25 time of from 1 minute to 20 hours will generally be suitable, in particular an incubation time of from 30 minutes to 10 hours will often be preferred.

The temperature of the reaction medium in the process of the invention may suitably be in the range of 10-90°C, in
30 particular in the range of 15-70°C, as appropriate for the XET enzyme in question.

The invention is further illustrated in the following non-limiting examples.

35

EXAMPLE 1

Screening for positive XET strains.

40 **Media**

- 5 PD Agar: 39 g potato dextrose agar, DIFCO 0013; add deionized water up to 1000 ml, autoclave at 121°C for 15 - 20 min.

YPG agar: 4 g yeast extract (DIFCO 0127),

1 g KH_2PO_4 (Merck 4873),

- 10 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (Merck 5886)

15 g dextrose (Roquette 101-0441)

20 g agar (Merck 1614)

deionized water up to 1000 ml

Autoclave at 121° C for 15 - 20 min.

- 15 MEA: 20 g malt extract powder (Difco 0186)

1 g peptone (Difco 0118)

20 g glucose (Roquette France 1010441)

20 g agar (Merck 1614)

deionized water up to 1000 ml

- 20 Autoclave at 121°C for 15 min

Medium A. Per flask: 30 g wheat bran , 45 ml of the following solution: 10 g rofec (Roquette 101-0441), 10 g NH_4NO_3 (Merck 1187), 10 g KH_2PO_4 (Merck 4873), 40 g Solcafloc (Dicacel available from Dicalite-Europe-Nord, 9000 Gent, Belgium), 0.75g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (Merck 5886), 15 g CaCO_3 , tap water to 1000 ml, pH adjusted to 6.5.

Autoclave for 40 min at 121° C.

- 5 Medium B. 20 g soyabean meal, 5 g maltodex 01, (Roquette 101-7845), 15 g wheat bran, 3 g peptone (Difco 0118), 10 g cellulose avicel (Merck 2331), 0.2 ml pluronic (PE-6100, 101-3068), 1 g olive oil, deionized water up to 1000 ml.
- 10 100 ml in 500 ml Erlenmeyer flask with 2 baffles. Autoclave at 121° C for 40 min.
- Medium C. 100 g sucrose, 40 g soybean meal, 10 g $\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$ (Merck 6579), 0.1 ml pluronic (PE 6100), tap water up to 1000 ml. 0.5 g CaCO_3 in a 2 baffled
- 15 Erlenmeyer flask with 100 ml medium. Autoclave at 121°C for 40 minutes. Just prior to use, add 10 ml of 1M NaHCO_3 pH 9 per 100 ml medium.

FERMENTATION PROCEDURE

- 20 The fungal strains were maintained on agar in petri dishes (9 cm) or on slants with PDA, YPG or MEA (see list of media). 1 agar slant was washed off with 10 ml sterile distilled water and 3 ml were used to inoculate 1 shake flask.

- The fungal strains were grown in shake flasks under the
- 25 following growth conditions:

media: A and B (see list of media)

Temperature: 26°C

RPM: A : stationary

B : 125 - 200

- 30 Incubation time : A : 6 to 30 days

B: 2 - 21 days.

- 5 The bacterial isolate was maintained on agar slopes with TY agar pH 9 which consists of:

20 g tryptone, 5 g yeast extract, 0.7 ml of a 1% solution of $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$, 0.1 ml of a 1% solution of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 1.5 ml of a 1% solution of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 20 g agar and deionized water up to 1000ml. Autoclave at 121°C for 20 minutes. Before pouring the plates, add 10 ml of sterile 1M NaHCO_3 pH9 to 100 ml agar.

The slants were incubated at 30°C for 4 days. 1 slant was washed off with 10 ml sterile distilled water and 3 ml were used to inoculate each flask.

- 15 The bacterium was grown in shake flasks containing Medium C at 30°C for 4 days at 300 rpm.

ACTIVITY TEST:

The culture broths which were produced in B and C, were centrifuged at 10,000 rpm for 10 minutes. The supernatants were tested for XET activity.

To each flask with wheat bran and fully grown culture, was added 200 ml tap water and shaken at 200 rpm for 2 hours. The culture broth was then centrifuged and the supernatant tested for XET.

25 XET ANALYSIS

The samples were analysed using the "dot-blot" assay described above.

RESULTS

Using the method described above, 1 *Bacillus* sp and several fungi belonging to different taxonomical groups were found to be positive. The results are shown in the table below.

The *Bacillus alcalophilus* was positive when grown on Medium C pH

5 9 (see list of media) for 4 days at 30°C at 300 rpm.

| Fungal Isolate | Medium A | Medium B |
|--------------------------------------|----------|----------|
| <i>Actinomucor elegans</i> | + | - |
| <i>Dichotomocladium hesseltinei</i> | + | + |
| <i>Gongronella butleri</i> | not done | + |
| <i>Mucor miehei</i> var <i>minor</i> | + | + |
| <i>Sporodiniella umbellata</i> | - | + |
| <i>Alternaria</i> sp | + | - |
| <i>Aposphaeria</i> | not done | + |
| <i>Aspergillus tamarii</i> | + | + |
| <i>Botrytis cinerea</i> | + | + |
| <i>Chaetapiozpora rhododendri</i> | - | + |
| <i>Colletotrichum acutatum</i> | + | - |
| <i>Colletotrichum crassipes</i> | + | - |
| <i>Coniothyrium dunkii</i> | + | + |
| <i>Coniothyrium olivaceum</i> | + | - |
| <i>Coniothyrium</i> sp | - | + |
| <i>Coryneum castaneicola</i> | - | + |
| <i>Cytospora</i> spp | + | + |
| <i>Diplodia gossypina</i> | + | + |
| <i>Discula</i> sp | + | + |
| <i>Embellisia hyacinthi</i> | + | - |
| <i>Eupenicillium javanicum</i> | + | - |

| | | |
|---------------------------------|----------|----------|
| <i>Eurotium chevalieri</i> | - | + |
| <i>Fusarium solani</i> | + | - |
| <i>Galiella celebica</i> | + | + |
| <i>Lulworthia uniseptata</i> | + | + |
| <i>Nodulisporium sp</i> | + | - |
| <i>Oedocephalum sp</i> | - | + |
| <i>Penicillium canescens</i> | + | - |
| <i>P. capsulatum</i> | + | - |
| <i>P. italicum</i> | not done | + |
| <i>P. olsonii</i> | + | - |
| <i>P. pinophilum</i> | + | not done |
| <i>P. roqueforti</i> | + | + |
| <i>P. verruculosum</i> | + | + |
| <i>Pestalotia sp</i> | not done | + |
| <i>Pestalotiopsis sp</i> | + | + |
| <i>Petromyces alliaceus</i> | + | + |
| <i>Phoma neoloba</i> | + | - |
| <i>Phoma tropica</i> | + | + |
| <i>Phomopsis cirsii</i> | - | + |
| <i>Phomopsis ilicis</i> | + | + |
| <i>Phyllosticta sp</i> | + | - |
| <i>Plowrightia ribesia</i> | + | + |
| <i>Poronia punctata</i> | + | + |
| <i>Pseudoplectania nigrella</i> | + | + |

| | | |
|-----------------------------------|---|----------|
| <i>Pyronema domesticum</i> | + | + |
| <i>Ramularia</i> sp | + | - |
| <i>Seimatosporium lichenicola</i> | + | - |
| <i>Septoria</i> sp | + | not done |
| <i>Talaromyces flavus</i> | + | + |
| <i>Tubeufia amazonensis</i> | + | - |
| <i>Tiarosporella phaseolina</i> | + | + |
| <i>Tiarosporella</i> sp | - | + |
| <i>Verticillium</i> sp | + | + |
| <i>Volutella buxi</i> | + | - |
| <i>Xylaria</i> sp | + | + |
| <i>Corticium roseum</i> | + | + |
| <i>Schizophyllum</i> sp | + | + |
| <i>Stereum hirsutum</i> | + | + |
| <i>Trametes hirsuta</i> | + | + |
| <i>Tubulicrinis subulatus</i> | + | + |
| <i>Acrodontium crateriforme</i> | + | - |
| <i>Aureobasidium pullulans</i> | + | - |
| <i>Circinotrichum</i> sp | - | + |
| <i>Cryptocline</i> sp | + | - |
| <i>Ellisiopsis</i> sp | + | - |
| <i>Epicoccum nigrum</i> | - | + |
| <i>Gliocladium</i> sp | + | - |
| <i>Helicorhoidion irregulare</i> | + | + |

| | | |
|-------------------------------|---|----------|
| <i>Hendersonia spp</i> | + | + |
| <i>Mariannaea sp</i> | + | - |
| <i>Microsphaeropsis sp</i> | + | - |
| <i>Ramularia sp</i> | + | - |
| <i>Sarcopodium sp</i> | + | - |
| <i>Spadicoides sp</i> | - | + |
| <i>Speiropsis pedatospora</i> | + | - |
| <i>Sporotrichum exile</i> | + | + |
| <i>Stilbella sp</i> | + | - |
| <i>Trichothecium sp</i> | + | + |
| <i>Trimmatostroma abietes</i> | + | + |
| <i>Tubakia dryina</i> | + | + |
| <i>Wiesneriomyces sp</i> | + | - |
| <i>Zygosporium masonii</i> | + | + |
| <i>Vialaea</i> | + | not done |

5 + = positive, - = negative.

EXAMPLE 2

Purification and characterization of *Dichotomocladium*

10 *hesseltinei* XET.

15 PDA agar slopes were inoculated with *Dichotomocladium hesseltinei* (CBS 164.61) and incubated at 26°C for 7 days. They were washed off with about 250 ml sterile distilled water with 0.1% Tween 80 and used to inoculate 80 shake flasks containing Medium B (2 - 3ml / flask). The flasks were shaken at 200 rpm,

- 5 26°C for 5 days after which time the culture broth was centrifuged at 4000 rpm for 15 minutes.

The supernatant containing xyloglucan endotransglucosylase (XET) was purified using the following method:

10

- Filter aid was added to the culture broth which was filtered through a filtration cloth. This solution was further filtered through a Seitz depth filter plate resulting in a clear solution. The pH of the filtrate was adjusted to pH 8.0 and the
15 filtrate was diluted with deionised water to give the same conductivity as 20mM Tris/HCl, pH 8.0.

- The XET enzyme was applied to a Q-sepharose FF column equilibrated in 20mM Tris/HCl, pH 8.0 and the enzyme was eluted
20 with an increasing linear NaCl gradient (0 → 0.5M). The XET activity eluted as a single peak. The pooled XET was transferred to 20mM Tris/HCl, pH 8.0 on a Sephadex G25 column and rechromatographed on a Q-sepharose FF column equilibrated in 20mM Tris/HCl, pH 8.0. The column was eluted with an increasing
25 linear NaCl gradient (0 → 0.2M). XET containing fractions were pooled and $(\text{NH}_4)_2\text{SO}_4$ was added to 1.4M final concentration. A Phenyl Toyopearl 650S column was equilibrated in 1.4M $(\text{NH}_4)_2\text{SO}_4$, 10mM succinic acid, pH 7.0 and the XET enzyme was applied to this column and eluted with a decreasing linear $(\text{NH}_4)_2\text{SO}_4$
30 gradient (1.4 → 0M). XET containing fractions were pooled and concentrated on an ultrafiltration cell with a 10kDa cut-off regenerated cellulose membrane. The concentrated enzyme was applied to a Superdex200 size exclusion column equilibrated in 100mM H_3BO_3 , 10mM dimethyl glutaric acid, 2mM CaCl_2 , pH 7.0.
35 Fractions eluted from the Superdex200 column were analyzed by SDS-PAGE and pure XET fractions were pooled.

- The *Dichotomocladium hesseltinei* XET migrates on SDS-PAGE as a band with $M_r = 24$ kDa. N-terminal amino acid sequencing of the
40 24 kDa component was carried out following SDS-PAGE and

- 5 electroblotting onto a PVDF-membrane. The following N-terminal amino acid sequence (SEQ ID No. 1) could be deduced:

Ala-Glu-Phe-Cys-Gly-Gln-Trp-Asp-Thr-Gln-Thr-Val-Gly-Asn-Tyr-Ile-
Val-Tyr-Asn-Asn-Leu-Leu-Gly-Ala-Gly-Ser-Ala.

10

- The present invention also relates to a microbial XET enzyme comprising the amino acid sequence shown in SEQ ID No. 1 or a XET being at least 80% homologous with the amino acid sequence SEQ ID No.1, preferably being at least 85% homologous with SEQ ID No. 1, more preferably being at least 90% homologous with SEQ ID No.1, even more preferably being at least 95% homologous with SEQ ID No. 1, in particular being at least 98% homologous with SEQ ID No. 1.

- A polypeptide is considered to be X% homologous to the parent XET if a comparison of the respective amino acid sequences, performed via known algorithms, such as the one described by Lipman and Pearson in Science 227, 1985, p. 1435, reveals an identity of X%.

- In addition, mass spec. analysis of the *Dichotomocladium hesseltinei* XET gave a value of 23 006 Da.

EXAMPLE 3

- 30 Purification and characterization of *Tiarosporella phaseolina* XET.

- 15 PDA agar slopes were inoculated with *Tiarosporella phaseolina* (CBS 446.97) and incubated at 26°C for 7 days. They were washed off with about 250 ml sterile distilled water with 0.1% Tween 80 and used to inoculate 80 shake flasks containing Medium B (2 - 3ml/flask). The flasks were shaken at 200 rpm, 26°C for 7 days after which time the culture broth was centrifuged at 4000 rpm for 15 minutes.

40

The supernatant containing xyloglucan endotransglucosylase (XET)

5 was purified using the following method:

Filter aid was added to the culture broth which was filtered through a filtration cloth. This solution was further filtered through a Seitz depth filter plate resulting in a clear
10 solution. The filtrate was concentrated by ultrafiltration on 3 kDa cut-off polyethersulfone membranes followed by dialfiltration with distilled water to reduce the conductivity. The pH of the concentrated enzyme was adjusted to pH 5.0. The conductivity of the concentrated enzyme was 1.7 mS/cm.

15 The XET enzyme was applied to a S-sepharose FF column equilibrated in 20mM CH₃COOH/NaOH, pH 5.0 and the enzyme was eluted with an increasing linear NaCl gradient (0 → 0.5M). The XET activity eluted as a single peak. The pooled XET was
20 transferred to 20mM CH₃COOH/NaOH, pH 4.0 by dialysis. The dialysed enzyme was applied to a SOURCE 30S column equilibrated in 20mM CH₃COOH/NaOH, pH 4.0. After washing the column the XET activity was eluted with an increasing linear NaCl gradient (0 → 0.5M). Fractions with XET activity were pooled and dialysed
25 against 20mM Tris/HCl, pH 8.0. The dialysed enzyme was applied to a SOURCE 30Q column equilibrated in 20mM Tris/HCl, pH 8.0. After washing the column the XET activity was eluted with an increasing linear NaCl gradient (0 → 0.5M). Fractions with XET activity were pooled and concentrated on an ultrafiltration cell
30 with a 10kDa cut-off regenerated cellulose membrane. The concentrated enzyme was applied to a Superdex200 size exclusion column equilibrated in 100mM H₃BO₃, 10mM dimethyl glutaric acid, 2mM CaCl₂, pH 7.0. Fractions eluted from the Superdex200 column were analyzed by SDS-PAGE and pure XET fractions were pooled.

35 The *Tiarosporrella phaseolina* XET migrates on SDS-PAGE as a band with $M_r = 24$ kDa. N-terminal amino acid sequencing of the 24 kDa component was carried out following SDS-PAGE and electroblotting onto a PVDF-membrane. The following N-terminal amino acid
40 sequence (SEQ. ID No. 2) could be deduced:

5

Xaa-Asp-Phe-Cys-Gly-Gln-Trp-Asp-Asn-Val-Lys-Asn-Gly-Pro-Tyr-Thr-
Leu-Tyr-Asn-Asn-Leu-Gly-Gly-Lys

The present invention also relates to a microbial XET enzyme
10 comprising the amino acid sequence shown in SEQ ID No. 2 or a XET
being at least 80% homologous with the amino acid sequence SEQ ID
No. 2, preferably being at least 85% homologous with SEQ ID No.
2, more preferably being at least 90% homologous with SEQ ID No.
2, even more preferably being at least 95% homologous with SEQ ID
15 No. 2, in particular being at least 98% homologous with SEQ ID
No. 2.

A polypeptide is considered to be X% homologous to the parent XET
if a comparison of the respective amino acid sequences, performed
via known algorithms, such as the one described by Lipman and
20 Pearson in Science 227, 1985, p. 1435, reveals an identity of X%.

EXAMPLE 4

Purification and characterization of *Pseudoplectania nigrella*

25 XET.

15 PDA agar slopes were inoculated with *Pseudoplectania nigrella*
(CBS 444.97) and incubated at 26°C for 7 days. They were washed
off with about 250 ml sterile distilled water with 0.1% Tween 80
30 and used to inoculated 80 shake flasks containing Medium B (2 -
3ml/flask). The flasks were shaken at 200 rpm, 26°C for 7 days
after which time the culture broth was centrifuged at 4000 rpm
for 15 minutes.

35 The supernatant containing xyloglucan endotransglycosylase (XET)
was purified using the following method:

Filter aid was added to the culture broth which was filtered
through a filtration cloth. This solution was further filtered
40 through a Seitz depth filter plate resulting in a clear
solution. The pH of the filtrate was adjusted to pH 5.0 and the

- 5 filtrate was diluted with deionised water to give the same conductivity as 20mM CH₃COOH/NaOH, pH 5.0.

The XET enzyme was applied to a S-sepharose FF column equilibrated in 20mM CH₃COOH/NaOH, pH 5.0 and the enzyme was
10 eluted with an increasing linear NaCl gradient (0 → 0.25M). The XET activity eluted as a single peak. The pooled XET was transferred to 20mM Hepes/NaOH, pH 7.0 on a Sephadex G25 column and applied to a Q-sepharose FF column equilibrated in 20mM Hepes/NaOH, pH 7.0. The column was eluted with an increasing
15 linear NaCl gradient (0 → 0.5M). XET containing fractions were pooled and transferred to 20mM CH₃COOH/NaOH, pH 5.0 by dialysis. A SOURCE 15S column was equilibrated in 20mM CH₃COOH/NaOH, pH 5.0, and the dialysed enzyme was applied. After washing the column the XET enzyme was eluted with an increasing linear NaCl
20 gradient (0 → 0.2M). XET containing fractions were pooled and concentrated on an ultrafiltration cell with a 10kDa cut-off regenerated cellulose membrane. The concentrated enzyme was applied to a Superdex200 size exclusion column equilibrated in 20mM CH₃COOH/NaOH, 100mM NaCl, pH 5.0. Fractions eluted from the
25 Superdex200 column were analyzed by SDS-PAGE and pure XET fractions were pooled.

The *Pseudoplectania nigrella* XET migrates on SDS-PAGE as a band with $M_r = 58$ kDa. Following SDS-PAGE and electroblotting onto a
30 PVDF-membrane it was found that the 58 kDa component had a blocked N-terminus.

A highly purified preparation of *Pseudoplectania nigrella* XET was reduced and alkylated. A sample of the enzyme was then
35 degraded with Lys-C. Peptides were isolated by RP-HPLC on a long Vydac C18 in a SMART system using TFA/AN and repurified in TFA/isopropanol. Selected peptides were analyzed by Edman degradation.

- 5 In Table 1 the sequences obtained from 8 peptides are shown. Two peptides were found to be homologous to glucanase- and xylanase-like enzymes but the majority did not show any or only irrelevant homology to known sequences.

10 Table 1. Sequence of Lys-C peptides from *P.nigrella* XET

| Peptide | Run no. | Sequence | Comments on homology |
|--------------------------|---------|---------------------|---|
| XET lysyl 070198-2fr5 | 3777 | WNDPVVK | Homology to B. subtilis lysis protein |
| XET lysyl 080198-4fr3 | 3778 | (S/Y)RFNAPALIGE[WQ] | Nothing found |
| XET lysyl 090198-1fr6 | 3779 | LIFE | Too small |
| XET lysyl 070198-1fr5 | P379 | EDGSYLKYAK | irrelevant homologies |
| XET lysyl 070198-1fr3 | P380 | EWGTTGKFNK | Homology to endoglucanase |
| XET lysyl 080198-1fr1 | P381 | KVTAVEAWK | Weak homology to xyn1_asptu. Endoxylanase |
| XET lysyl 080198-2fr1 | P382 | XFYQIANS(Q/I) | Nothing found |
| XET lysyl 080198-3fr1 | P383 | AALXXVMK | Nothing found |

15 **EXAMPLE 5**

pH profile of *Dichotomocladium hesseltinei*, *Tiarosporella phaseolina* and *Pseudoplectania nigrella* XET and xyloglucanase

- 20 The xyloglucan endotransglycosylase of *Dichotomocladium hesseltinei*, *Tiarosporella phaseolina* and *Pseudoplectania*

5 *nigrella* were checked for their pH profile. The pure enzymes were diluted in buffers ranging from pH 3.0 to 11.0 so that in the final dilution, the protein concentration was the same i.e. $A_{280} = 0.004$. The samples were then assayed for XET using the XET-dot-blot-assay (described earlier). The fluorescent spots
10 were judged visually and graded from 0 - 10. The units used in Fig. 1 are therefore arbitrary units.

It can be seen from Fig. 1 that the XET is active between pH 3 and pH 11, in particular between pH 4 and pH 9.

15 All the isolates with XET activity were also tested for xyloglucanase activity. The ratio between the 2 enzymes varies from isolate to isolate. The enzymes were diluted in the same buffers as for the XET pH profile activity until the final protein concentration was the same and assayed for xyloglucanase
20 activity. AZCL xyloglucan was used. The xyloglucanase method used was the following:

Substrate: 0.4% AZCL-xyloglucan suspended in demineralised water.

25 Buffer: 100mM H_3BO_3 , 10mM Dimethy glutaric acid, 2mM $CaCl_2$, pH 7.

Analysis:

- An Eppendorf thermomixer is switched on at 40° C
- 500µl 0.4% AZCL-xyloglucan is mixed with 500µl buffer and put
30 on ice.
- 20µl enzyme is added and incubated in the Eppendorf thermomixer until a suitable colour is reached.
- The samples are returned to the ice bath to prevent further reaction while the samples are centrifuged at 0°C.
- 35 - 200 µl samples are transferred to micro titre plates and the blue colour is measured at 650nm.

The results are shown in Fig. 2.

5 **EXAMPLE 6**

Cloning and expression of a xyloglucan endotransglycosylase enzyme (XET) from *Tiarosporella phaseolina* CBS No. 446.97

10 Deposited organisms:

Tiarosporella phaseolina CBS No. 446.97.

Other strains:

Yeast strain: The *Saccharomyces cerevisiae* strain used was W3124

- 15 (van den Hazel, H.B; Kielland-Brandt, M.C.; Winther, J.R. in Eur. J. Biochem., 207, 277-283, 1992; (MATa; ura 3-52; leu 2-3, 112; his 3-D200; pep 4-1137; prc1::HIS3; prb1::LEU2; cir+).

E. coli strain: DH10B (Life Technologies)

Plasmids:

- 20 The *Aspergillus* expression vector pHD414 is a derivative of the plasmid p775 (described in EP 238 023). The construction of pHD414 is further described in WO 93/11249.

pYES 2.0 (Invitrogen)

Media:

- 25 **YPD:**

10 g yeast extract, 20 g peptone, H₂O to 900 ml. Autoclaved, 100 ml 20% glucose (sterile filtered) added.

YPM:

- 10 g yeast extract, 20 g peptone, H₂O to 900 ml. Autoclaved, 100
30 ml 20% maltodextrin (sterile filtered) added.

10 x Basal salt:

75 g yeast nitrogen base, 113 g succinic acid, 68 g NaOH, H₂O ad 1000 ml, sterile filtered.

SC-URA:

- 35 100 ml 10 x Basal salt, 28 ml 20% casamino acids without vitamins, 10 ml 1% tryptophan, H₂O ad 900 ml, autoclaved, 3.6 ml 5% threonine and 100 ml 20% glucose or 20% galactose added.

SC-agar:

SC-URA, 20 g/l agar added.

- 5 AZCL Xyloglucan (Megazyme, Australia)

Expression cloning in yeast

Expression cloning in yeast was done as described by H. Dalboege et al. (H. Dalboege et al Mol. Gen. Genet (1994) 243:253-260.; WO 93/11249; WO 94/14953), which are hereby incorporated as
10 reference. All individual steps of Extraction of total RNA, cDNA synthesis, Mung bean nuclease treatment, Blunt-ending with T4 DNA polymerase, and Construction of libraries was done according to the references mentioned above.

Fermentation procedure of *Tiarosporella phaseolina* CBS No.

- 15 446.97 for mRNA isolation:

Tiarosporella phaseolina CBS No. 446.97 was inoculated from a plate with outgrown mycelium into a shake flask containing 100 ml medium B (see media). The culture was incubated at 26°C and 200 rpm for 7 days. The resulting culture broth was filtered
20 through miracloth and the mycelium was frozen down in liquid nitrogen.

mRNA was isolated from mycelium from this culture as described in (H. Dalboege et al Mol. Gen. Genet (1994) 243:253-260.; WO 93/11249; WO 94/14953).

- 25 Extraction of total RNA was performed with guanidinium thiocyanate followed by ultracentrifugation through a 5.7 M CsCl cushion, and isolation of poly(A)⁺RNA was carried out by oligo(dT)-cellulose affinity chromatography using the procedures described in WO 94/14953.

- 30 cDNA synthesis:

Double-stranded cDNA was synthesized from 5 mg poly(A)⁺ RNA by the RNase H method (Gubler and Hoffman (1983) Gene 25:263-269, Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY). The poly(A)⁺
35 RNA (5 µg in 5 µl of DEPC-treated water) was heated at 70°C for 8 min. in a pre-siliconized, RNase-free Eppendorph tube, quenched on ice and combined in a final volume of 50 µl with reverse transcriptase buffer (50 mM Tris-Cl, pH 8.3, 75 mM KCl, 3 mM MgCl₂, 10 mM DTT, Bethesda Research Laboratories)
40 containing 1 mM of dATP, dGTP and dTTP and 0.5 mM 5-methyl-dCTP

5 (Pharmacia), 40 units human placental ribonuclease inhibitor (RNasin, Promega), 1.45 μ g of oligo(dT)₁₈-Not I primer (Pharmacia) and 1000 units SuperScript II RNase H reverse transcriptase (Bethesda Research Laboratories). First-strand cDNA was synthesized by incubating the reaction mixture at 45°C
10 for 1 hour. After synthesis, the mRNA:cDNA hybrid mixture was gelfiltrated through a MicroSpin S-400 HR (Pharmacia) spin column according to the manufacturer's instructions.

After the gelfiltration, the hybrids were diluted in 250 μ l second strand buffer (20 mM Tris-Cl, pH 7.4, 90 mM KCl, 4.6 mM
15 MgCl₂, 10 mM (NH₄)₂SO₄, 0.16 mM bNAD+) containing 200 μ l of each dNTP, 60 units *E. coli* DNA polymerase I (Pharmacia), 5.25 units RNase H (Promega) and 15 units *E. coli* DNA ligase (Boehringer Mannheim). Second strand cDNA synthesis was performed by incubating the reaction tube at 16°C for 2 hours and additional
20 15 min. at 25°C. The reaction was stopped by addition of EDTA to a final concentration of 20 mM followed by phenol and chloroform extractions.

Mung bean nuclease treatment:

The double-stranded cDNA was precipitated at -20°C for 12 hours
25 by addition of 2 vols 96% EtOH, 0.2 vol 10 M NH₄Ac, recovered by centrifugation, washed in 70% EtOH, dried and resuspended in 30 μ l Mung bean nuclease buffer (30 mM NaAc, pH 4.6, 300 mM NaCl, 1 mM ZnSO₄, 0.35 mM DTT, 2% glycerol) containing 25 units Mung bean nuclease (Pharmacia). The single-stranded hair-pin DNA was
30 clipped by incubating the reaction at 30°C for 30 min., followed by addition of 70 μ l 10 mM Tris-Cl, pH 7.5, 1 mM EDTA, phenol extraction and precipitation with 2 vols of 96% EtOH and 0.1 vol 3 M NaAc, pH 5.2 on ice for 30 min.

Blunt-ending with T4 DNA polymerase:

35 The double-stranded cDNAs were recovered by centrifugation and blunt-ended in 30 ml T4 DNA polymerase buffer (20 mM Tris-acetate, pH 7.9, 10 mM MgAc, 50 mM KAc, 1 mM DTT) containing 0.5 mM of each dNTP and 5 units T4 DNA polymerase (New England Biolabs) by incubating the reaction mixture at 16°C for 1 hour.
40 The reaction was stopped by addition of EDTA to a final

5 concentration of 20 mM, followed by phenol and chloroform extractions, and precipitation for 12 hours at -20°C by adding 2 vols 96% EtOH and 0.1 vol 3 M NaAc pH 5.2.

Adaptor ligation, Not I digestion and size selection:

After the fill-in reaction the cDNAs were recovered by
10 centrifugation, washed in 70% EtOH and dried. The cDNA pellet was resuspended in 25 µl ligation buffer (30 mM Tris-Cl, pH 7.8, 10 mM MgCl₂, 10 mM DTT, 0.5 mM ATP) containing 2.5 µg non-palindromic BstXI adaptors (Invitrogen) and 30 units T4 ligase (Promega) and incubated at 16°C for 12 hours. The reaction was
15 stopped by heating at 65°C for 20 min. and then cooling on ice for 5 min. The adapted cDNA was digested with Not I restriction enzyme by addition of 20 µl water, 5 µl 10x Not I restriction enzyme buffer (New England Biolabs) and 50 units Not I (New England Biolabs), followed by incubation for 2.5 hours at 37°C.
20 The reaction was stopped by heating at 65°C for 10 min. The cDNAs were size-fractionated by gel electrophoresis on a 0.8% SeaPlaque GTG low melting temperature agarose gel (FMC) in 1x TBE to separate unligated adaptors and small cDNAs. The cDNAs were size-selected with a cut-off at 0.7 kb and rescued from the
25 gel by use of b-Agarase (New England Biolabs) according to the manufacturer's instructions and precipitated for 12 hours at -20°C by adding 2 vols 96% EtOH and 0.1 vol 3 M NaAc pH 5.2.

Construction of library:

The directional, size-selected cDNAs were recovered by
30 centrifugation, washed in 70% EtOH, dried and resuspended in 30 µl 10 mM Tris-Cl, pH 7.5, 1 mM EDTA. The cDNAs were desalted by gel filtration through a MicroSpin S-300 HR (Pharmacia) spin column according to the manufacturer's instructions. Three test ligations were carried out in 10 µl ligation buffer (30 mM Tris-
35 Cl, pH 7.8, 10 mM MgCl₂, 10 mM DTT, 0.5 mM ATP) containing 5 µl double-stranded cDNA (reaction tubes #1 and #2), 15 units T4 ligase (Promega) and 30 ng (tube #1), 40 ng (tube #2) and 40 ng (tube #3, the vector background control) of BstXI-NotI cleaved pYES 2.0 vector. The ligation reactions were performed by
40 incubation at 16°C for 12 hours, heating at 70°C for 20 min. and

5 addition of 10 μ l water to each tube. 1 μ l of each ligation mixture was electroporated into 40 μ l electrocompetent *E. coli* DH10B cells (Bethesda research Laboratories) as described (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY). Using the
10 optimal conditions a library was established in *E. coli* consisting of pools. Each pool was made by spreading transformed *E. coli* on LB+ampicillin agar plates giving 15.000-30.000 colonies/plate after incubation at 37°C for 24 hours. 20 ml LB+ampicillin was added to the plate and the cells were
15 suspended herein. The cell suspension was shaken in a 50 ml tube for 1 hour at 37°C. Plasmid DNA was isolated from the cells according to the manufacturer's instructions using QIAGEN plasmid kit and stored at -20°C.

1 μ l aliquots of purified plasmid DNA (100 ng/ml) from
20 individual pools were transformed into *S. cerevisiae* W3124 by electroporation (Becker and Guarante (1991) Methods Enzymol. 194:182-187) and the transformants were plated on SC agar containing 2% glucose and incubated at 30°C.

Identification of positive colonies:

25 Colonies were screened indirectly for XET by finding xyloglucanase positive colonies.

After 3-5 days of growth, the agar plates were replica plated onto a set of SC-URA agar (with 20% galactose added) plates containing 0.1% AZCL Xyloglucan. These plates were incubated for
30 3-7 days at 30°C. Xyloglucanase positive colonies were identified as colonies surrounded by a blue halo.

Cells from enzyme-positive colonies were spread for single colony isolation on agar, and an enzyme-producing single colony was selected for each of the Xyloglucanase-producing colonies identified.
35

All xyloglucanase positive colonies were tested for XET and were found to be positive.

Isolation of a cDNA gene for expression in *Aspergillus*:

An XET-producing yeast colony was inoculated into 20 ml YPD
40 broth in a 50 ml glass test tube. The tube was shaken for 2 days

5 at 30°C. The cells were harvested by centrifugation for 10 min.
at 3000 rpm.

DNA was isolated according to WO 94/14953 and dissolved in 50 ml
water. The DNA was transformed into *E. coli* by standard
procedures. Plasmid DNA was isolated from *E. coli* using standard
10 procedures, and analyzed by restriction enzyme analysis. The
cDNA insert was excised using the restriction enzymes BamH I and
Xba I and ligated into the *Aspergillus* expression vector pHD414
resulting in the plasmid pA2XG80.

The cDNA inset of Qiagen purified plasmid DNA of pA2XG80
15 (Qiagen, USA) was sequenced with the Taq deoxy terminal cycle
sequencing kit (Perkin Elmer, USA) and synthetic oligonucleotide
primers using an Applied Biosystems ABI PRISM™ 377 DNA
Sequencer according to the manufacturers instructions.

Transformation of *Aspergillus oryzae* or *Aspergillus niger*

20 Protoplasts are prepared as described in WO 95/02043, p. 16,
line 21 - page 17, line 12, which is hereby incorporated by
reference.

100 µl of protoplast suspension is mixed with 5-25 µg of the
appropriate DNA in 10 µl of STC (1.2 M sorbitol, 10 mM Tris-HCl,
25 pH = 7.5, 10 mM CaCl₂). Protoplasts are mixed with p3SR2 (an *A.*
nidulans amdS gene carrying plasmid) (Tove Christensen et al.
Bio/Technology, pp 1419-1422 vol.6, Dec. 1988). The mixture is
left at room temperature for 25 minutes. 0.2 ml of 60% PEG 4000
(BDH 29576), 10 mM CaCl₂ and 10 mM Tris-HCl, pH 7.5 is added and
30 carefully mixed (twice) and finally 0.85 ml of the same solution
is added and carefully mixed. The mixture is left at room
temperature for 25 minutes, spun at 2500 g for 15 minutes and
the pellet is resuspended in 2 ml of 1.2 M sorbitol. After one
more sedimentation the protoplasts are spread on minimal plates
35 (Cove, Biochem. Biophys. Acta 113 (1966) 51-56) containing 1.0 M
sucrose, pH 7.0, 10 mM acetamide as nitrogen source and 20 mM
CsCl to inhibit background growth. After incubation for 4-7 days
at 37°C spores are picked and spread for single colonies. This
procedure is repeated and spores of a single colony after the

- 5 second reisolation is stored as a defined transformant.

Test of *A. oryzae* transformants

Each of the *A. oryzae* transformants is inoculated in 10 ml of YPM (cf. below) and propagated. After 2-5 days of incubation at 30°C, the supernatant is removed.

- 10 The XET activity is identified by using the XET dot-blot assay described earlier.

The following sequence (SEQ ID No. 3) is the cDNA insert in pYES2.0 of XG80 (XET from *Tiarosporella phaseolina*) including the BamH I and Not I recognition sites used for cloning:

- 15 GGATCCGAATTCCAACATATCCTGCCCTCCTTTCAAGCGAACACCATGAAGTTCTCCTCGGCTCT
GTTTCTGGCCGCTACG
GCGGTCTTGGCTTCCGCCGCGCCGCTTGAGCGCCGCGCCGACTTTTGTGGTCAATGGGACAACG
TGAAGAACGGACCTTA
20 CACTCTTTACAACAACCTGTGGGAAAAGATGCTTCCGGAGCCTCCGGATCGCAATGCACCGGC
GTCGATAGCTTCAGCA
GCAACACCATCGCTTGGCACACATCCTGGTCCTGGTCCGGTGCTCAGTACAATGTCAAGTCTTA
CGCAAACGTGGTTCGTC
GACATCACCTCTAAGAAACTCAGCGCCATCAGCAGCATTAAACAGCATCTGGCGCTGGGCTTACA
25 CGGGTAGCAACATTGT
TGCCAATGTTGCCTACGATATCTTCACTTCATCCACTGTCGGTGGTAGCGAGGAATATGAAATC
ATGATATGGGTTGGTG
CTCTCGGTGGTGCTGGTCCGATCTCATCTACCGGCTCCCCTATTGCCACCGTTTCCCTTGCAGG
CTCCTCGTGGAAGCTC
30 TACAAAGGGCCCAACGGGCAGATGACCGTGTTTCACTTCGTCGCGGAGTCCAACGTGAACAACT
TCAGCGGTGACCTTAA
CGCTTTCATCAAGTACCTCACCGCAACCAGGGCCTTCCCGCCTCGCAATACATCAAGAGCATT
GGCGCTGGCACTGAGC
CGTTACGGGTTCACACGCCAAGCTGACCACTTCCTCCTACACTGTCAGCTTCAGATAACTGTG
35 AAGCTTTATGCTGCCC
TTATGCATCATCCTTGTTACATAGTTATCACCAGGGGACTCTTGTAATACGATTGCCTTATTAA
CCGCCTGCATCTGCTT
TCACACAATGGCATTACCAATCAACAGTGCGCCTCGAATCCGTAAAAGGTGGCTTAAAAAAA
AAAAAAA
40 AATTCCTGCGGCCGC.

The following sequence (SEQ ID no. 4) is the amino acid sequence of the coding region of XG80:

- 45 MKFSSALFLAATAVLASAAPLERRADFCGQWDNVKNGPYTLNNLWGKDASGASGSQCTGVDSF
SSNTIAWHTSWSWSGA
QYNVKSANVVVDITSKKLSAIISSINSIWRWAYTGSNIVANVAYDIFTSSTVGGSEYEIMIWV
GALGGAGPISSTGSP
ATVSLAGSSWKLYKGPNGQMTVFSFVAESNVNNSGDLNFIKYLGNQGLPASQYIKSIGAGT
50 EPFTGSNAKLTTSSYT
VSFR.

- 5 The present invention also relates to a microbial XET enzyme comprising the amino acid sequence shown in SEQ ID No. 4 or a XET being at least 80% homologous with the amino acid sequence SEQ ID No. 4, preferably being at least 85% homologous with SEQ ID No. 4, more preferably being at least 90% homologous with SEQ ID No. 4, even more preferably being at least 95% homologous with SEQ ID No. 4, in particular being at least 98% homologous with SEQ ID No. 4.

- 10 A polypeptide is considered to be X% homologous to the parent XET if a comparison of the respective amino acid sequences, performed via known algorithms, such as the one described by Lipman and Pearson in Science 227, 1985, p. 1435, reveals an identity of X%.

A clone Cl.XG80 was deposited at DSMZ on 24 February 1998 under Accession No. DSM 12032.

5 SEQUENCE LISTING

(1) GENERAL INFORMATION:

10 (i) APPLICANT:

(A) NAME: NOVO NORDISK A/S
(B) STREET: Novo Alle
(C) CITY: Bagsvaerd
(E) COUNTRY: Denmark
15 (F) POSTAL CODE (ZIP): DK-2880
(G) TELEPHONE: +45 44 44 88 88
(H) TELEFAX: +45 44 49 32 56

20 (ii) TITLE OF INVENTION: MICROBIAL XYLOGLUCAN ENDOTRANSGLYCOSYLASE
(XET)

(iii) NUMBER OF SEQUENCES: 4

25 (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

30 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 27 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: peptide

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ala Glu Phe Cys Gly Gln Trp Asp Thr Gln Thr Val Gly Asn Tyr Ile
1 5 10 15
Val Tyr Asn Asn Leu Leu Gly Ala Gly Ser Ala
20 25

50 (2) INFORMATION FOR SEQ ID NO: 2:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

60 (ii) MOLECULE TYPE: peptide

65 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Xaa Asp Phe Cys Gly Gln Trp Asp Asn Val Lys Asn Gly Pro Tyr Thr
1 5 10 15
Leu Tyr Asn Asn Leu Gly Gly Lys

5

20

(2) INFORMATION FOR SEQ ID NO: 3:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 975 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: DNA (genomic)

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

| | |
|--|-----|
| GGATCCGAAT TCCTAATATC CTGCCCCTCT TTCAAGCGAA CACCATGAAG TTCTCCTCGG | 60 |
| CTCTGTTTCT GGCCGCTACG GCGGTCTTGG CTTCCGCCGC GCCGCTTGAG CGCCGCGCCG | 120 |
| ACTTTTGTGG TCAATGGGAC AACGTGAAGA ACGGACCTTA CACTCTTTAC AACAACTGT | 180 |
| GGGGAAAAGA TGCTTCCGGA GCCTCCGGAT CGCAATGCAC CGGCGTCGAT AGCTTCAGCA | 240 |
| GCAACACCAT CGCTTGGCAC ACATCCTGGT CCTGGTCCGG TGCTCAGTAC AATGTCAAGT | 300 |
| CTTACGCAAA CGTGGTCGTC GACATCACCT CTAAGAACT CAGCGCCATC AGCAGCATT | 360 |
| ACAGCATCTG GCGCTGGGCT TACACGGGTA GCAACATTGT TGCCAATGTT GCCTACGATA | 420 |
| TCTTCACTTC ATCCACTGTC GGTGGTAGCG AGGAATATGA AATCATGATA TGGGTTGGTG | 480 |
| CTCTCGGTGG TGCTGGTCCG ATCTCATCTA CCGGCTCCCC TATTGCCACC GTTTCCTTG | 540 |
| CAGGCTCCTC GTGGAAGCTC TACAAAGGGC CCAACGGGCA GATGACCGTG TTCAGCTTCG | 600 |
| TCGCCGAGTC CAACGTGAAC AACTTCAGCG GTGACCTTAA CGCTTTCATC AAGTACCTCA | 660 |
| CCGGCAACCA GGGCCTTCCC GCCTCGCAAT ACATCAAGAG CATTGGCGCT GGCACGAGC | 720 |
| CGTTCACGGG TTCCAACGCC AAGCTGACCA CTTCTCCTA CACTGTCAGC TTCAGATAAC | 780 |
| TGTGAAGCTT TATGCTGCCC TTATGCATCA TCCTTGTTACA TAGTTATCAC CAGGGGACTC | 840 |
| TTGTAAATAC GATTGCCTTA TTAACCGCCT GCATCTGCTT TCACACAATG GCATTTACCA | 900 |
| ATCAACAGTG CGCCTCGAAT CCGTAAAAGG TGGCTTAAAA AAAAAAAAAA AAAAAAAAAA | 960 |
| AATTCCTGCG GCCGC | 975 |

55

(2) INFORMATION FOR SEQ ID NO: 4:

60

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 244 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

65

(ii) MOLECULE TYPE: peptide

70

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

5 Met Lys Phe Ser Ser Ala Leu Phe Leu Ala Ala Thr Ala Val Leu Ala
 1 5 10 15
 Ser Ala Ala Pro Leu Glu Arg Arg Ala Asp Phe Cys Gly Gln Trp Asp
 20 25 30
 10 Asn Val Lys Asn Gly Pro Tyr Thr Leu Tyr Asn Asn Leu Trp Gly Lys
 35 40 45
 15 Asp Ala Ser Gly Ala Ser Gly Ser Gln Cys Thr Gly Val Asp Ser Phe
 50 55 60
 Ser Ser Asn Thr Ile Ala Trp His Thr Ser Trp Ser Trp Ser Gly Ala
 65 70 75 80
 20 Gln Tyr Asn Val Lys Ser Tyr Ala Asn Val Val Val Asp Ile Thr Ser
 85 90 95
 Lys Lys Leu Ser Ala Ile Ser Ser Ile Asn Ser Ile Trp Arg Trp Ala
 100 105 110
 25 Tyr Thr Gly Ser Asn Ile Val Ala Asn Val Ala Tyr Asp Ile Phe Thr
 115 120 125
 30 Ser Ser Thr Val Gly Gly Ser Glu Glu Tyr Glu Ile Met Ile Trp Val
 130 135 140
 Gly Ala Leu Gly Gly Ala Gly Pro Ile Ser Ser Thr Gly Ser Pro Ile
 145 150 155 160
 35 Ala Thr Val Ser Leu Ala Gly Ser Ser Trp Lys Leu Tyr Lys Gly Pro
 165 170 175
 Asn Gly Gln Met Thr Val Phe Ser Phe Val Ala Glu Ser Asn Val Asn
 180 185 190
 40 Asn Phe Ser Gly Asp Leu Asn Ala Phe Ile Lys Tyr Leu Thr Gly Asn
 195 200 205
 45 Gln Gly Leu Pro Ala Ser Gln Tyr Ile Lys Ser Ile Gly Ala Gly Thr
 210 215 220
 Glu Pro Phe Thr Gly Ser Asn Ala Lys Leu Thr Thr Ser Ser Tyr Thr
 225 230 235 240
 50 Val Ser Phe Arg

5


| | | | |
|---|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|---|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|---|--------------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page 9, line 17-22 | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | |
| Further deposits are identified on an <input checked="" type="checkbox"/> | |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 448.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is <input type="checkbox"/> continued on an additional sheet | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5


| | | |
|---|---------|--|
| Applicant's or agent's file reference number | 5154-WO | International application No. PCT/DK 98/00076 |
|---|---------|--|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|--|---|
| A. The indications made below relate to the microorganism referred to in the description on page 9, line 28 to page 10, line 5 | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 2 January 1996 | Accession Number CBS 831.95 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is <input type="checkbox"/> continued on an additional sheet Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5


| | | | |
|---|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|---|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

| | |
|---|---|
| A. The indications made below relate to the microorganism referred to in the description on page 10, lines 6-12 | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 12 March 1996 | Accession Number CBS 274.96 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is <input type="checkbox"/> continued on an additional sheet | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

10

| | |
|---|--|
| For receiving Office use only | For International Bureau use only |
| <input checked="" type="checkbox"/> This sheet was received with the international application | <input type="checkbox"/> This sheet was received by the International Bureau on: |
| Authorized officer  | Authorized officer |

5


| | | | |
|---|---------|------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No | PCT/DK 98/00076 |
|---|---------|------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

10

(PCT Rule 13bis)

| | |
|--|-------------------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page 11, lines 7-13 | |
| B. IDENTIFICATION OF DEPOSIT | |
| Further deposits are identified on an additional sheet | <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 446.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| |
|---|
| For receiving Office use only |
| <input checked="" type="checkbox"/> This sheet was received with the international application |
| Authorized officer  |

| |
|--|
| For International Bureau use only |
| <input type="checkbox"/> This sheet was received by the International Bureau on: |
| Authorized officer |

5

| | | | |
|---|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|---|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

10

(PCT Rule 13bis)

| | |
|---|---|
| <p>A. The indications made below relate to the microorganism referred to in the description on page 11, lines 20-26</p> | |
| <p>B. IDENTIFICATION OF DEPOSIT additional sheet</p> | <p>Further deposits are identified on an <input checked="" type="checkbox"/> additional sheet</p> |
| <p>Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES</p> | |
| <p>Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands</p> | |
| <p>Date of deposit 28 January 1997</p> | <p>Accession Number CBS 444.97</p> |
| <p>C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/></p> <p>Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited.</p> | |
| <p>D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)</p> | |
| <p>E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)</p> <p>The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")</p> | |
| <p>For receiving Office use <input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer <i>[Signature]</i></p> | <p>For International Bureau use <input type="checkbox"/> This sheet was received by the International Bureau only on:</p> <p>Authorized officer</p> |

5


| | | | |
|---|---------|------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No | PCT/DK 98/00076 |
|---|---------|------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|--|---|
| A. The indications made below relate to the microorganism referred to in the description on page 11, line 31 to page 12, line 7 | |
| B. IDENTIFICATION OF DEPOSIT | |
| additional sheet | Further deposits are identified on an <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 447.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is <input type="checkbox"/> continued on an additional sheet Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |
| | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5

| | | | |
|---|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|---|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|---|-------------------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page 12, lines 10-16 | |
| B. IDENTIFICATION OF DEPOSIT | |
| Further deposits are identified on an additional sheet | <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 445.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|--|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer <i>E. van der Linden</i> | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5


| | | | |
|---|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|---|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

10

(PCT Rule 13bis)

| | |
|---|---|
| A. The indications made below relate to the microorganism referred to in the description on page 12, line 30 to page 13, line 6. | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input checked="" type="checkbox"/> X |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 2 January 1996 | Accession Number CBS 830.95 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5

| | | | |
|---|---------|-------------------------------|-------------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98 / 00076 |
|---|---------|-------------------------------|-------------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

| | |
|--|--|
| A. The indications made below relate to the microorganism referred to in the description on page 13, lines 12-18. | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input checked="checked" type="checkbox"/> X |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 442.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> | |
| <p>Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited.</p> | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

For receiving Office use
only

☒ This sheet was received with the international
application

Authorized officer

Ernest J. Fitch

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

5

| | | | |
|--|---------|-------------------------------|-----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98/00076 |
|--|---------|-------------------------------|-----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

10

(PCT Rule 13bis)

| | |
|---|---|
| A. The indications made below relate to the microorganism referred to in the description on page 13, lines 23-29. | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input checked="" type="checkbox"/> X |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 23 January 1997 | Accession Number CBS 424.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| |
|--|
| For receiving Office use only |
| <input checked="" type="checkbox"/> This sheet was received with the international application |
| Authorized officer <i>Ernst J. J. J. J.</i> |

| |
|--|
| For International Bureau use only |
| <input type="checkbox"/> This sheet was received by the International Bureau on: |
| Authorized officer |

5


| | | | |
|---|---------|-------------------------------|-------------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK 98 / 00076 |
|---|---------|-------------------------------|-------------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|--|-------------------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page 13, line 30 to page 14, line 6. | |
| B. IDENTIFICATION OF DEPOSIT | |
| Further deposits are identified on an additional sheet | <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 23 January 1997 | Accession Number CBS 425.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5


| | | | |
|---|---------|-------------------------------|----------------|
| Applicant's or agent's file reference number | 5154-WO | International application No. | PCT/DK98/00076 |
|---|---------|-------------------------------|----------------|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|--|-------------------------------------|
| A. The indications made below relate to the microorganism referred to in the description on page 15, line 29 to page 16, line 6. | |
| B. IDENTIFICATION OF DEPOSIT | |
| Further deposits are identified on an additional sheet | <input checked="" type="checkbox"/> |
| Name of depositary institution CENTRAALBUREAU VOOR SCHIMMELCULTURES | |
| Address of depositary institution (including postal code and country) Oosterstraat 1, Postbus 273, NL-3740 AG Baarn, The Netherlands | |
| Date of deposit 28 January 1997 | Accession Number CBS 443.97 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| | |
|---|--|
| For receiving Office use only | |
| <input checked="" type="checkbox"/> | This sheet was received with the international application |
| Authorized officer  | |

| | |
|-----------------------------------|---|
| For International Bureau use only | |
| <input type="checkbox"/> | This sheet was received by the International Bureau on: |
| Authorized officer | |

5

| | |
|---|---|
| Applicant's or agent's file reference number 5154-WO | International application No. PCT/DK 98/00076 |
|---|---|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|---|--|
| A. The indications made below relate to the microorganism referred to in the description on page 17, lines 11-16. | |
| B. IDENTIFICATION OF DEPOSIT | |
| additional sheet | Further deposits are identified on an <input checked="checked" type="checkbox"/> |
| Name of depositary institution DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELLKULTUREN GmbH | |
| Address of depositary institution (including postal code and country) Mascheroder Weg 1b, D-38124 Braunschweig, GERMANY | |
| Date of deposit 12 February 1997 | Accession Number DSM 11404 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is <input type="checkbox"/> | |
| continued on an additional sheet | |
| Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

For receiving Office use
only

☒ This sheet was received with the international application

Authorized officer

James H. [Signature]

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

5


| | |
|---|---|
| Applicant's or agent's file reference number 5154-WO | International application No. PCT/DK 98/00076 |
|---|---|

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

10

| | |
|--|--|
| A. The indications made below relate to the microorganism referred to in the description on page 45, lines 17-18. | |
| B. IDENTIFICATION OF DEPOSIT additional sheet | Further deposits are identified on an <input type="checkbox"/> |
| Name of depositary institution DEUTSCHE SAMMLUNG VON MIKROORGANISMEN UND ZELLKULTUREN GmbH | |
| Address of depositary institution (including postal code and country) Mascheroder Weg 1b, D-38124 Braunschweig, GERMANY | |
| Date of deposit 24 February 1998 | Accession Number DSM 12032 |
| C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/> Until the publication of the mention of grant of a European patent or, where applicable, for twenty years from the date of filing if the application has been refused, withdrawn or deemed withdrawn, a sample of the deposited microorganism is only to be provided to an independent expert nominated by the person requesting the sample (cf. Rule 28(4) EPC). And as far as Australia is concerned, the expert option is likewise requested, reference being had to Regulation 3.25 of Australia Statutory Rules 1991 No 71. Also, for Canada we request that only an independent expert nominated by the Commissioner is authorized to have access to a sample of the microorganism deposited. | |
| D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) | |
| E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) | |
| The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") | |

| |
|---|
| For receiving Office use only |
| <input checked="" type="checkbox"/> This sheet was received with the international application |
| Authorized officer  |

| |
|--|
| For International Bureau use only |
| <input type="checkbox"/> This sheet was received by the International Bureau on: |
| Authorized officer |

5 CLAIMS

1. A method for the production of a xyloglucan endotransglycosylase enzyme (XET) comprising
 - (a) culturing in a suitable nutrient medium a microorganism
 - 10 expressing a microbial XET under conditions conducive to the production of the XET enzyme, and
 - (b) subsequently recovering of the XET enzyme from the nutrient medium.
- 15 2. The method according to claim 1, wherein the microorganism is a fungus or a bacterium.
3. The method according to claim 2, wherein the fungus is a basidiomycota, an ascomycota, a zygomycota or a mitosporic
- 20 fungus.
4. The method according to claim 3, wherein the fungus is a basidiomycotinum strain of the order *Coriolales*, *Schizophyllales*, *Stereales*, or *Xenasmatales*.
- 25 5. The method according to claim 4, wherein the fungus is a basidiomycotinum selected from a strain belonging to the group consisting of the families *Coriolaceae*, *Corticaceae*, *Schizophyllaceae*, *Stereaceae* and *Tubulicrinaceae*.
- 30 6. The method according to claim 4, wherein the fungus is a basidiomycotinum selected from a strain belonging to the group consisting of the genera *Trametes*, *Corticium*, *Schizophyllum*, *Stereum* and *Tubulicrinis*.
- 35 7. The method according to claim 6, wherein the fungus is a basidiomycotinum selected from a strain belonging to the group consisting of the species *Trametes hirsuta*, *Corticium roseum*, *Schizophyllum sp*, *Stereum hirsutum* and *Tubulicrinis subulatus*.

5 8. The method according to claim 7, wherein the fungus is a *Schizophyllum sp.*, deposit no. CBS 443.97.

9. The method according to claim 3, wherein the ascomycetes is selected from a strain belonging to the group consisting of the
10 classes *Loculoascomycetes*, *Discomycetes*, *Pyrenomycetes*, and *Plectomycetes*.

10. The method according to claim 9, wherein the ascomycetes is selected from a strain belonging to the group consisting of the
15 orders *Dothideales*, *Rhytismatales*, *Pezizales*, *Leotiales*, *Xylariales*, *Hypocreales*, *Halosphaeriales*, *Eurotiales*, *Phyllachorales* and *Diaporthales*.

11. The method according to claim 10, wherein the ascomycetes is
20 selected from a strain belonging to the group consisting of the families *Leptosphaeriaceae*, *Botryosphaeriaceae*, *Dothioraceae*, *Mycosphaerellaceae*, *Tubeufiaceae*, *Rhytismataceae*, *Sarcosomataceae*, *Pyronemataceae*, *Sclerotiniaceae*, *Amphisphaeriaceae*, *Hyponectriaceae*, *Xylariaceae*, *Valsaceae*,
25 *Melanconidaceae*, *Hypocreaceae*, *Halosphaeriaceae*, *Phyllachoraceae* and *Trichocommataceae*.

12. The method according to claim 11, wherein the ascomycetes is selected from a strain belonging to the group consisting of the
30 genera *Coniothyrium*, *Phoma*, *Diplodia*, *Plowrightia*, *Phyllosticta*, *Septoria*, *Tubeufia*, *Alternaria*, *Embellisia*, *Tiarosporella*, *Galiella*, *Oedocephalum*, *Pseudoplectania*, *Pyronema*, *Botrytis*, *Aposphaeria*, *Pestalotia*, *Pestalotiopsis*, *Chaetapiozpora*, *Poronia*, *Nodulisporium*, *Xylaria*, *Cytospora*, *Discula*, *Phomopsis*, *Coryneum*,
35 *Seimatosporium*, *Fusarium*, *Verticillium*, *Volutella*, *Lulworthia*, *Colletotrichum*, *Aspergillus*, *Eurotium*, *Eupenicillium*, *Penicillium*, *Petromyces* and *Talaromyces*.

13. The method according to claim 12, wherein the ascomycetes is

5 selected from a strain belonging to the group consisting of the
species *Diplodia gossypina*, *Plowrightia ribesia*, *Phyllosticta* sp,
Septoria sp, *Tubeufia amazonensis*, *Alternaria* sp, *Embellisia*
hyacinthi, *Phoma neoloba*, *Phoma tropica*, *Coniothyrium* sp,
Coniothyrium olivaceum, *Coniothyrium dunckii*, *Tiarosporella* sp,
10 *Tiarosporella phaseolina*, *Galiella celebica*, *Pseudoplectania*
nigrella, *Pyronema domesticum*, *Oedocephalum* sp, *Botrytis cinerea*,
Aposphaeria sp, *Pestalotia* sp, *Pestalotiopsis* sp, *Poronia*
punctata, *Xylaria* sp, *Nodulisporium* sp, *Fusarium solani*,
Verticillium sp, *Volutella buxi*, *Chaetapiozpora rhododendri*,
15 *Lulworthia uniseptata*, *Colletotrichum aculatum*, *Colletotrichum*
crassipes, *Cytospora* spp, *Discula* sp, *Phomopsis* sp, *Phomopsis*
cirsii, *Coryneum castaneicola*, *Seimatosporium lichenicola*,
Aspergillus tamaris, *Eurotium chevalieri*, *Eupenicillium*
javanicum, *Penicillium capsulatum*, *Penicillium olsonii*,
20 *Penicillium pinophilum*, *Penicillium roqueforti*, *Penicillium*
italicum, *Penicillium verruculosum*, *Penicillium canescens*,
Petromyces alliaceus and *Talaromyces flavus*.

14. The method according to claim 13, wherein the ascomycetes is
25 selected from a strain belonging to the group consisting of the
species *Botrytis cinerea* deposit no. CBS 447.97, *Pseudoplectania*
nigrella deposit no. CBS 444.97, *Tiarosporella phaseolina* deposit
no. CBS 446.97, *Pestalotia* sp deposit no. CBS 445.97 and
Lulworthia uniseptata deposit no. CBS 442.97.

30 15. The method according to claim 3, wherein the zygomycota is
selected from a strain belonging to the order Mucorales.

16. The method according to claim 15, wherein the fungus is a zy-
35 gomycotum selected from a strain belonging to the group
consisting of the families *Chaetocladiaceae* and *Mucoraceae*.

17. The method according to claim 16, wherein the fungus is a zy-

5 gomycotum selected from a strain belonging to the group
consisting of the genera *Dichotomocladium*, *Actinomucor*,
Gongronella, *Sporodiniella*, and *Mucor*.

18. The method according to claim 17, wherein the fungus is a zy-
10 gomycota selected from a strain belonging to the group consisting
of the species *Dichotomocladium hesseltinei*, *Actinomucor*
elegans, *Gongronella butleri*, *Mucor miehei* var *minor* and
Sporodiniella umbellata.

15 19. The method according to claim 18, wherein the fungus is a
Gongronella butleri deposit no. CBS 448.97.

20. The method according to claim 2, wherein the fungus is a
mitosporic fungus.

20

21. The method according to claim 1, wherein the microorganism is
Vialaea insculpta.

22. The method according to claim 2, wherein the bacterium is
25 gram-positive.

23. The method according to claim 22, wherein the gram-positive
bacterium is *Bacillus*.

30 24. The method according to claim 23, wherein the gram-positive
bacterium is *Bacillus alcalophilus* deposit no DSM 11404.

25. Use of a XET preparation obtained according to any of claims
1-24 for treating a cellulosic material.

35

26. The use according to claim 25, wherein the treatment provides
improved strength and/or improved shape-retention and/or improved
anti-wrinkling properties of the cellulosic material.

5 27. The use according to claim 25, wherein the cellulosic material is a cellulosic fabric or a paper and pulp product.

28. A xyloglucan endotransglycosylase preparation which is producible by cultivation of a microorganism expressing a microbial
10 XET.

29. The preparation according to claim 28, wherein the XET has the amino acid sequence shown in SEQ ID No. 1, or the XET has an amino acid sequence which is at least 80% homologous with SEQ ID
15 No. 1.

30. The preparation according to claim 28, wherein the XET has the amino acid sequence shown in SEQ ID No. 2, or the XET has an amino acid sequence which is at least 80% homologous with SEQ ID
20 No. 2.

31. The preparation according to claim 28, wherein the XET has the amino acid sequence shown in SEQ ID No. 4, or the XET has an amino acid sequence which is at least 80% homologous with SEQ ID
25 No. 4.

32. The preparation according to claim 28, wherein the XET is active between 3 and 11, in particular between 4 and 9.

30 33. A method for the production of a xyloglucan endotransglycosylase enzyme (XET) comprising
(a) culturing in a suitable nutrient medium a transformed host microorganism expressing a microbial XET under conditions conducive to the production of the XET enzyme, and
35 (b) subsequently recovering of the XET enzyme from the nutrient medium.

1/2

XET units




Fig. 1

2/2

XYLOGLUCANASE UNITS




Fig. 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00076

| | | |
|---|--|--|
| A. CLASSIFICATION OF SUBJECT MATTER | | |
| IPC6: C12N 9/10, C12N 9/24, C12N 9/42 According to International Patent Classification (IPC) or to both national classification and IPC | | |
| B. FIELDS SEARCHED | | |
| Minimum documentation searched (classification system followed by classification symbols) | | |
| IPC6: C12N | | |
| Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched | | |
| SE,DK,FI,NO classes as above | | |
| Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) | | |
| WPI, CA, MEDLINE, BIOSIS | | |
| C. DOCUMENTS CONSIDERED TO BE RELEVANT | | |
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| A | WO 9513384 A1 (UNILEVER PLC), 18 May 1995 (18.05.95) | 1-24 |
| X | -- | 25-33 |
| A | EP 0562836 A1 (TAKARA SHUZO CO.LTD.), 29 Sept 1993 (29.09.93) | 1-24 |
| X | -- | 25-33 |
| <input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex. | | |
| * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family | | |
| Date of the actual completion of the international search | | Date of mailing of the international search report |
| 25 May 1998 | | 17 -06- 1998 |
| Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86 | | Authorized officer Yvonne Siösteen Telephone No. +46 8 782 25 00 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00076

| C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT | | |
|---|---|-----------------------|
| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
| A | Dialog Information Service, File Biosis, Dialog accession no. 7153746, Biosis accession no. 93138746, Fry S. C. et al: "Xyloglucan Endotransglycosylase A New Wall -Loosening Enzyme Activity From Plants", Biochem J 282 (3). 1992. 821-828 | 1-24 |
| X | <p style="text-align: center;">-- -----</p> | 25-33 |

INTERNATIONAL SEARCH REPORT
Information on patent family members

29/04/98

International application No.
PCT/DK 98/00076

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9513384 A1 | 18/05/95 | AU 8112994 A | 29/05/95 |
| | | CA 2176133 A | 18/05/95 |
| | | CZ 9601361 A | 11/12/96 |
| | | EP 0728208 A | 28/08/96 |
| | | HU 74598 A | 28/01/97 |
| | | HU 9601232 D | 00/00/00 |
| | | JP 9511121 T | 11/11/97 |
| | | PL 317046 A | 03/03/97 |
| EP 0562836 A1 | 29/09/93 | SK 57696 A | 01/10/96 |
| | | AU 667706 B | 04/04/96 |
| | | AU 3540593 A | 30/09/93 |
| | | CA 2092366 A | 27/09/93 |
| | | JP 6086670 A | 29/03/94 |
| | | JP 7079778 A | 28/03/95 |
| | | US 5516694 A | 14/05/96 |